

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
6 January 2005 (06.01.2005)

PCT

(10) International Publication Number
WO 2005/001496 A1

(51) International Patent Classification⁷: **G01R 33/28**

(21) International Application Number:
PCT/GR2004/000037

(22) International Filing Date: 29 June 2004 (29.06.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
20030100282 30 June 2003 (30.06.2003) GR

KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).


(71) Applicant and
(72) Inventor: PAPPAS, Panagiotis, T. [GR/GR]; 26, Markopoulioti Street, GR-11744 Athens (GR).

Published:
— with international search report
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHOD AND MEANS OF MULTI-ACTIVATION OF IONS AND ATOMS WITH NMR AND EPR.



8

(57) Abstract: The invention uses one inductor, which is comprised of one, two or few twisted or parallel conductors and exposes the sample in a pulsed and damped alternating magnetic field B without necessarily the employment of a second constant intensity magnetic field. In this way, the nuclei and the electrons of the sample-object are activated, in the presence of a non-constant magnetic field B that gets infinite negative and positive values between successive damped positive and negative values, crossing through the zero value during a magnetic pulse. Thus, a wide nuclear NMR and Electronic EPR Multiple - Resonance of sample, is achieved.

WO 2005/001496 A1

BEST AVAILABLE COPY